

ENEE 661, Nonlinear Control Systems

PSK

02-21-00

Lecture 4 (part II)

One of the basic results of single variable calculus in the classical mean value theorem (MVT)

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ . There is  $c, a < c < b$  such that the derivative

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

The adjoining picture gives us an idea of what's going on. The essential geometric idea is that




Figure 1 (MVT)

the tangent to the graph of  $f$  is parallel to the line joining points  $(a, f(a))$  and  $(b, f(b))$ .

Let us see what happens in higher dimensions. Consider  $f: [a, b] \rightarrow \mathbb{R}^2$




Figure 2. (not MVT)

If the curve defined by  $f$  is "convex" then, there is no  $c$  such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

the tangent to the curve is parallel to the line joining  $(a, f(a))$  and  $(b, f(b))$  in the  $x-y-z$  space. The classical MVT does not hold for a specific

Example:  $f: [0, \frac{\pi}{2}] \rightarrow \mathbb{R}^2$

$$x \mapsto (\cos(x), \sin(x))$$

$$f(b) - f(a) = (-1, 1) \in \mathbb{R}^2$$

$$b - a = \frac{\pi}{2}$$

There does not exist  $c \in [0, \frac{\pi}{2}]$  such that  $\frac{1}{\frac{\pi}{2}} \cdot (-\sin(c), \cos(c)) = (-1, 1)$  since it

$$\text{would require } \sin^2(c) + \cos^2(c) = \frac{11}{8} \neq 1.$$

The correct form of mean value theorem in higher dimension is actually an inequality. We need some preliminary results.

Lemma 1

Let  $a < b$   $f: [a, b] \rightarrow V$  a normed linear space and  $g: [a, b] \rightarrow \mathbb{R}$ ,  $f$  and  $g$  continuous on  $[a, b]$  and differentiable on  $(a, b)$ . Suppose,

$$\|f'(t)\| \leq g'(t) \quad a < t < b.$$

$$\text{Then } \|f(b) - f(a)\| \leq g(b) - g(a).$$

PSK  
02-21-00

Proof (of Lemma 1):  $\|f(b) - f(a)\| = \left\| \int_a^b f'(t) dt \right\|$

$$\leq \int_a^b \|f'(t)\| dt$$

$$\leq \int_a^b g(t) dt$$

$$= g(b) - g(a)$$

□

Lemma 2

Same hypothesis as in Lemma 1, except that the condition on existence and inequality of derivative holds for all  $t \in [a, b]$  except for a countable set of points. Same conclusion as Lemma 1.

Proof: (essentially same argument as in Lemma 1 hence the integrals are unaffected)

Corollary 3

Some hypothesis on  $f$  as in Lemma 1, and  $g(t) = kt, k > 0$ . Then  $\|f'(t)\| \leq k \forall t \in (a, b)$ .

Then,

$$\|f(b) - f(a)\| \leq k(b-a)$$

□

We need the definition of derivative for maps.

Definition Let  $E, F$  be normed linear spaces over  $\mathbb{R}$ . Let  $U \subset \mathbb{R}$  open. Suppose

$$f: U \rightarrow F$$

We say  $f$  is differentiable at  $a \in U$  if there is a continuous linear map

$$L: E \rightarrow F$$

such that

$$\lim_{h \rightarrow 0} \|f(a+h) - f(a) - Lh\|_F = 0,$$

$$h \rightarrow 0$$

$$\|h\|_E$$

here  $h$  is such that  $h+a \in U$ .

Clearly, if  $L$  exists it is unique,

and is given by

$$L(k) = \lim_{t \rightarrow 0} \frac{f(a+kt) - f(a)}{t}$$

$$= \left. \frac{d}{dt} f(a+kt) \right|_{t=0}$$

We call  $L$  the derivative of  $f$  at

$a$  and denote it by  $(Df)_a$  and

sometimes by  $Df(a)$ .

Exercise: (a) Chain rule

$$D(g \circ f)(a) = Dg(f(a)) \circ Df(a)$$

Composition of

nonlinear maps

of linear maps.

$$(b) \quad E = \mathbb{R}^n \text{ \& } F = \mathbb{R}^m$$

$$(Df)(a) \cdot h = \begin{bmatrix} a^T f'' h \\ \text{grad } f \end{bmatrix}$$

Jacobian

Mean Value Theorem 4

Let  $f: U \subset E \rightarrow F$  be a map of normed linear spaces. Let  $[a, b]$  denote the line segment

$$\{(1-t)a + tb : 0 \leq t \leq 1\},$$

with end-points  $a, b \in U$ , contained in  $U$ .

Then,

$$\|f(b) - f(a)\| \leq \sup_{t \in [0,1]} \|Df[(1-t)a + tb]\| \cdot \|b-a\|$$



Proof: Simply restrict  $f$  to the line segment  $[a, b]$  and then apply Corollary 3 above

Another useful result from calculus is

The Fundamental Theorem of Integral Calculus

Let  $X$  and  $Y$  be Banach spaces. Let  $U \subset E$  be an open set and  $f: U \rightarrow Y$  be a differentiable, everywhere in  $U$ , map in  $C^1$  map. Suppose  $a + ty \in U \forall t \in [0, 1]$ . Then

$$f(x+ty) = f(x) + \int_0^1 Df(x+ty)y dt$$

PSK  
62.21.0

Proof: The completeness / Banach property is used in the proper definition of integral with all attendant properties, as in 1 variable calculation. We take this for granted. Then, set  $g(t) = f(x+ty)$  for  $0 < t < 1$  by chain rule.

$$g'(t) = \frac{d}{dt} f(x+ty) y$$

$$\text{Let } k(t) = f(x) + \int_0^t \frac{d}{ds} f(x+sy) y ds$$

$$\text{Then } k'(t) = \frac{d}{dt} f(x+ty) y \quad 0 < t < 1$$

$$\text{Hence } g'(t) = k'(t)$$

$$\Rightarrow g(t) = k(t) + \text{constant} \quad 0 < t < 1$$

By continuity of  $g$  &  $h$  (they are integrals)  
 $g(t) = k(t) + \text{constant} \quad 0 \leq t \leq 1$   
 But  $g(0) = k(0) = f(x)$

$$\text{So } g(t) = f(x+ty) \quad \square$$

Lemma (Differentiability and Lipschitz Condition).

Let  $f: [a, b] \times D \rightarrow \mathbb{R}^n$  for domain  $D \subset \mathbb{R}^n$

continuous in  $t$  and  $(\frac{\partial f}{\partial x})$  exists and is continuous on  $[a, b] \times D$ . Then  $f$  is locally

Lipschitz on  $[a, b] \times \mathbb{D}$ .

Proof: For  $x_0 \in \mathbb{D}$ , let  $r > 0$  be such that

$$\mathbb{D}_0 \equiv \{x \mid \|x - x_0\| \leq r\} \subset \mathbb{D}.$$

$\mathbb{D}_0$  is closed and bounded.  $\mathbb{D}_0$  is convex,

since, for  $x_1, x_2 \in \mathbb{D}_0$  and  $0 \leq \alpha \leq 1$ ,

$$\|\alpha x_1 + (1-\alpha)x_2 - x_0\| = \|\alpha x_1 + (1-\alpha)x_2 - \alpha x_0 - (1-\alpha)x_0\|$$

$$= \|\alpha(x_1 - x_0) + (1-\alpha)(x_2 - x_0)\|$$

$$\leq \alpha \|x_1 - x_0\| + (1-\alpha) \|x_2 - x_0\|$$

$$\leq \alpha r + (1-\alpha)r$$

$$= r.$$

Then, by the Mean Value Theorem,  $\forall x, y \in \mathbb{D}_0$

$$\|f(t, y) - f(t, x)\|$$

$$\leq \sup_{0 \leq s \leq 1} \|Df((1-s)x + sy)\| \cdot \|y - x\|$$

$$\leq \sup_{t \in [a, b]} \sup_{0 \leq s \leq 1} \|Df((1-s)x + sy)\| \cdot \|y - x\|$$

$$= L \cdot \|y - x\|,$$

where we used continuity wrt both  $t$  &  $x$  of  $Df$  in the sup norms.

address: (02/26/02)

Suppose  $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is  $C^1$  (i.e. continuously differentiable) at each point  $x$  of an open set  $S \subset \mathbb{R}^n$ .

Suppose  $x^*, y^* \in S$  one such that the line

segment  $L(x^*, y^*)$  joining  $x^*$  and  $y^*$  is  $S$ . Then

there exists a point  $\bar{x} \in L(x^*, y^*)$  such that

$$f(y^*) - f(x^*) = \left( \frac{\partial f}{\partial x} \right)_{x=\bar{x}} (y^* - x^*)$$

Proof let  $g(t) = f((1-t)x^* + ty^*)$

then  $g(0) = f(x^*)$

$g(1) = f(y^*)$

$$g'(t) = \left( \frac{\partial f}{\partial x} \right)_{x=(1-t)x^* + ty^*} \cdot \frac{d}{dt} ((1-t)x^* + ty^*)$$

$$= \left( \frac{\partial f}{\partial x} \right)_{x=\bar{x}} \cdot (y^* - x^*)$$

By the scalar mean value theorem  $(g(1) - g(0)) = g'(t) \cdot (1-0)$

which is the same as saying

$$f(y^*) - f(x^*) = \left( \frac{\partial f}{\partial x} \right)_{x=\bar{x}} \cdot (y^* - x^*)$$

where  $\bar{x} = (1-t)x^* + ty^* \in L(x^*, y^*)$

